

James Madison University

## JMU Scholarly Commons

---

Doctor of Nursing Practice (DNP) Final Clinical  
Projects

The Graduate School

---

Fall 2019

### The effect of provider education on discharge information grade level and readability

Amber V. Balzer

Follow this and additional works at: <https://commons.lib.jmu.edu/dnp201019>



Part of the [Nursing Commons](#)

---

#### Recommended Citation

Balzer, Amber V., "The effect of provider education on discharge information grade level and readability" (2019). *Doctor of Nursing Practice (DNP) Final Clinical Projects*. 34.  
<https://commons.lib.jmu.edu/dnp201019/34>

This Dissertation is brought to you for free and open access by the The Graduate School at JMU Scholarly Commons. It has been accepted for inclusion in Doctor of Nursing Practice (DNP) Final Clinical Projects by an authorized administrator of JMU Scholarly Commons. For more information, please contact [dc\\_admin@jmu.edu](mailto:dc_admin@jmu.edu).

The Effect of Provider Education on  
Discharge Information Grade Level and Readability

Amber V. Balzer

A Clinical Research Project  
submitted to the Graduate Faculty of

JAMES MADISON UNIVERSITY

In

Partial Fulfillment of the Requirements

for the degree of

Doctor of Nursing Practice

School of Nursing

December 2019

---

FACULTY COMMITTEE:

Committee Chair: Erica Lewis RN, PhD

Committee Members/ Readers:

Dr. Rehan Qayyum MBBS, MHS, SFHM, FAHA

## Dedication & Acknowledgments

I dedicate this work to my husband, Phillip, who has been my biggest supporter in the pursuit of this degree. I would also like to acknowledge the faculty and advisors who have provided unconditional support and enthusiasm throughout this project.

## Table of Contents

Dedication and Acknowledgements.....	ii
List of Tables.....	v
Abstract.....	vi
I.    Introduction.....	1
II.   Problem Description.....	1
Background and Significance.....	2
Rationale.....	7
Purpose.....	8
III.  Methods.....	9
Location.....	9
Contextual Elements.....	10
Ethical Considerations.....	10
Baseline Measures.....	11
Intervention.....	11
Team.....	13
Outcome Measures.....	13
CPD-Reaction Questionnaire.....	13
Provider Self-Report.....	15
Post-Intervention DIF Scores.....	15
Data Collection Methods.....	15
Analysis.....	16
CPD-Reaction Questionnaire.....	16

	Provider Self-Report.....	18
	Post-Intervention DIF Scores.....	19
	Results.....	19
	Demographic Information.....	19
	CPD-Reaction Questionnaire.....	20
	Provider Self Report.....	22
	Pre/Post-Intervention DIF Scores.....	23
IV.	Discussion.....	24
	CPD-Reaction Questionnaire.....	25
	Provider Self-Report.....	27
	Post-Intervention DIF Scores.....	27
	Limitations.....	30
	Conclusion.....	30
V.	Appendices.....	32
	PRIDE Survey.....	32
	The Four Level Model of the Healthcare System.....	35
	Research Announcement.....	36
	Poster Presentation.....	37
	CPD-Reaction Questionnaire.....	38
	Self-Report Survey.....	41
VI.	References.....	42

## **List of Tables**

Table 1. Construct Calculations

Table 2. Demographic Information

Table 3. Construct Comparison

Table 4. Paired Samples Test

Table 5. Mann-Whitney Ranking

Table 6. Mann-Whitney Testing

## Abstract

**Problem:** The peri-discharge period for patients can be a complex and dangerous time. A large urban academic medical center with a readmission rate approaching 16% utilizes a standardized template called a Discharge Information Form (DIF) to convey information to inpatients at the time of their discharge. The problem is a lack of individualization with a universal template such as the DIF. Moreover, the readability and literacy level of the DIF have not been previously assessed.

**Methods:** A random sample of DIFs was assessed for literacy level using the Flesch-Kincaid Readability Test Tool. An in-person educational intervention was administered to a sample of Hospital Medicine Advanced Practice Providers with the objective of improving their understanding of DIF readability. The providers completed the Continuing Professional Development-Reaction Questionnaire before and after the educational intervention to assess clinical behavioral intentions. One-month post intervention a survey was administered to assess self-reported behavior change, and descriptive statistics were used to look for trends in these data. DIFs were also reassessed for readability, and a paired sample t-test was performed to determine if a change occurred.

**Findings:** After the intervention, statistically significant differences were found in the construct of *social influence* ( $p=0.040$ ). Grade level of DIFs was significantly improved ( $p=0.001$ ), readability of DIFs was nearly significantly improved ( $p=0.051$ ), and the majority of providers self-reported behavior change.

**Conclusion:** Providers responsible for creating patient discharge information should know how to write literacy level appropriate materials. There is an opportunity to improve discharge information readability through inpatient provider education.

## **Introduction**

In the acute care setting, all patients are given written discharge instructions at the time of their release from the hospital. Information conveyed in these instructions includes crucial topics such as medications, summary of the hospital stay, specific disease process instructions, activity restrictions, dietary restrictions, symptoms requiring calls to 911 or their primary care provider, and follow-up appointments. In order for patients to follow these instructions and remain out of the hospital, the instructions must be written in a way that patients are able to comprehend (Institute for Healthcare Improvement, 2019).

A priority of the Affordable Care Act is to reduce hospital readmissions (NEJM Catalyst, 2018). To meet this initiative, Medicare instituted the Hospital Readmissions Reduction Program (HRRP) which penalizes acute-care hospitals with 30-day readmission rates higher than similar health care systems (NEJM Catalyst, 2018). With evidence demonstrating that patients with low health literacy are more likely to experience hospital admissions and longer lengths of stay (Choudhry et al., 2015), training providers on techniques to improve both readability and grade level of written materials is both an ethical and financial responsibility of health systems.

## **Problem Description**

A large urban academic medical center has a 30-day readmission rate approaching 16%. The Division of Hospital Medicine within this center discharged 6,149 patients in 2018. This constitutes over 19% of the total number of patients discharged from the institution. Current practice for this division is to utilize a standardized template called a Discharge Information Form (DIF) to convey information to all inpatients at the time of their discharge from the hospital. With a universal template such as the DIF, individualization to patient needs can be lacking, which is problematic. Moreover, the readability and literacy level of the DIF have not



been previously assessed. In a time when patient-centered care and patient satisfaction are increasingly critical to the delivery of healthcare, it is vital to educate the providers responsible for creating the DIF on the individualized needs of patients based on their reported preferences.

In order to have an understanding of patient preferences within the large urban academic medical center, the Patient Perceptions of Relative Importance of Discharge Elements (PRIDE) Study was designed. The PRIDE study was a cross-sectional survey completed in early 2019. The survey (see Appendix A) allowed patients to rank the relative importance of common elements of discharge instructions and collected basic demographic information. Results of this study determined that regardless of age, race, gender, educational level, and medical comorbidities, patients admitted to this hospital perceived medication information as having the highest relative importance (Balzer et al., 2019).

### **Background and Significance**

The peri-discharge period can be a complex and dangerous time for patients discharged from the local health system, with nearly 16% of patients requiring readmission. When considering ways to prevent the dangers associated with this period, the Agency for Healthcare Research and Quality (2018) suggests that structured and patient-centered discharge communication is crucial. The Joint Commission mandates information that must be included in all discharge communication including; the reason for hospitalization, significant findings, procedures and treatment provided, patient's discharge condition, patient and family instructions, and attending physician signature (Horwitz et al., 2013). This study focused on the area of patient and family instructions, which typically includes topics such as discharge medications, activity orders, therapy orders, dietary instructions, and medical follow up. In previous studies at

the national level, readability, understandability, and completeness of these areas have been shown to be subpar (Unaka et al., 2017).

Health literacy refers to a patient's ability "to obtain, process, and understand basic health information and services needed to make appropriate health decisions" (U.S. Department of Health and Human Services, 2010). Health literacy affects numerous aspects of hospitalization including navigating the healthcare system, completing complex forms, locating providers and services, sharing personal information, engaging in self-care and chronic-disease management, and understanding mathematical concepts, such as probability and risk (Health Literacy Basics, 2017). Factors such as stress and sickness have been shown to decrease health literacy, and lower health literacy rates are linked to increased hospitalizations as well as readmissions (Agency for Healthcare Research and Quality, 2013). Therefore, the Joint Commission's (2007) accreditation standards highlight the need for patients to receive information about their care in a way in which they can understand.

In order to be effectively understood by the average adult, patient and family instructions should be written at a 6th-grade level, according to the National Institutes of Health, the US Department of Health and Human Services (2000), and the American Medical Association (1999) (as cited in Choudhry et al, 2016). Despite these guidelines, discharge instructions are frequently written at a much higher level, often requiring a college graduate reading level to comprehend (Choudhry et al., 2016). Providers should focus on strategies to increase awareness in matching patient-oriented information to appropriate levels of literacy (Choudhry et al., 2016), and in order to accomplish this task, providers must both be aware of the educational levels of their patients, and have an understanding of the preferences associated with each educational level.

A variety of other factors affect how patients comprehend discharge instructions including age, sex, medical comorbidities, and language. It has been found that increasing age has been associated with non-comprehension of medications and follow up instructions, male sex has been associated with non-comprehension of diet instructions, and depression has been associated with medication non-compliance (Albrecht et al., 2014). 8.6% of the United States population demonstrates Limited English Proficiency and are 40% more likely to experience physical harm from an adverse event than their English-speaking counterparts (Agency for Healthcare Research and Quality, 2018). In order to provide quality care, it is crucial for providers to understand how a variety of factors influence the way patients receive and interpret discharge communication.

Efforts to improve discharge information readability nationwide have demonstrated some success. Boston University Medical Center and the Agency for Healthcare Research and Quality created the Re-Engineered Discharge (RED) which has proven effective at reducing readmissions and posthospital emergency department visits, while also improving patient satisfaction by enforcing twelve components of the discharge process (Boston University, 2014). These components include “identify the correct medicines and a plan for the patient to obtain and take them, teach a written discharge plan the patient can understand, educate the patient about his or her diagnosis, review with the patient what to do if a problem arises” (Boston University, 2014). Many of these identified components fall into the category of patient and family instructions which will be the focus of this study. The Joint Commission also supports efforts made by health systems to meet the Standards of Care for Transitions, which is a set of guidelines created by the multistakeholder Transition of Care Consensus Conference (as cited in, Polster, 2015).

On a local level, Marcantoni, Finney, & Lane (2015) aimed to improve the literacy level of discharge communication by utilizing an educational brochure written at a seventh-grade reading level compared to traditional discharge teaching and found that the standardized lower literacy level tool improved post-hospital follow-up adherence. Mueller, Giannelli, Boxer, & Schnipper (2015) found that the use of disease specific templated discharge instructions was associated with better readability than the use of clinician-generated discharge instructions. Waniga, Gerke, Shoemaker, Bourgoine, & Eamranond (2016) found that patient satisfaction in the discharge domain was significantly increased after the implementation of a nurse and physician generated standardized teaching tool targeting discharge communication.

Numerous studies have been completed with the goal of improving the hospital discharge process. El-Eid, Kaddoum, Tamim, & Hitti (2015) found that improving the timeliness of hospital discharge by aiming to have patients discharged before noon each day can decrease length of stay, save money, and improve hospital and emergency department throughput. Kwan, Bell, Morgan, & Stewart (2013) found that implementing a patient navigator within the healthcare team to coordinate patient care, communicate with patients and families, and oversee care transitions can decrease length of stay. Some studies aimed at increasing support after discharge through interventions such as a discharge phone call allowing staff to provide better, more detailed discharge instructions to patients after they arrived home gave improved medication management, follow-up appointment reminders, and opportunities for answering patient questions (Schuller, Lin, Gamm, & Edwardson, 2015). Efforts to improve care transitions perceived as the most helpful by patients include speaking to the pharmacist, receiving an illustrated medication schedule, receiving a follow-up phone call at home, and getting help from friends and family (Cawthon et al., 2013).

Some studies have focused on improving hospital discharge through the nursing staff providing discharge instructions. One of the most important duties nurses have is to prepare patients for discharge by teaching patients about their conditions, medications, self-care strategies and the importance of follow-up care (Jakucs, 2018). One method explored educating nurses to utilize the teach-back method to ensure patients understood discharge instructions, which demonstrated an increase in patient satisfaction (Burke, 2018). Another method encouraged utilizing “do not disturb” signs to create an uninterrupted block of time for discharge teaching (Polster, 2015). Despite the large amounts of literature surrounding the discharge process, few studies have focused on directly educating the providers responsible for creating discharge instructions.

Gaps in existing literature have been identified including limited data suggesting how healthcare providers believe discharge communication should be prioritized. Blaine, et al. (2018) indicated discharge education/teach-back and involving the care team as areas perceived by providers as having the highest importance, and Sorita, et al. (2017) notes medical history, physical findings, cognitive and functional status at discharge, and rationale for medication changes to be “very important.” Although patient satisfaction with discharge information strongly correlates to overall satisfaction with hospital care (Waniga, Gerke, Shoemaker, Bourgoine, & Eamranond 2016), there is limited literature in which patients rate their perceived relative importance of discharge information, and no evidence deciphering whether educational level plays a role in relative importance rankings. Corser, Dontje, Neuberger, Chant, & Keskimaki (2017) found that 44% of patients felt that improvements were needed in the areas of formatting/layout, clarity, correcting discrepancies/omitted information, or providing the document in electronic form, therefore determining how receiving this information is prioritized

and improving readability could help to make discharge communication more patient-centered to prevent readmissions and adverse events.

Given the lack of information about provider prioritization of discharge instructions nationally, and the general lack of information within the local health system about current readability of discharge information, this study was needed. The general purpose of this study was to assess the readability of local discharge information, present providers with practical tools to improve the readability of written instructions, and determine if a change in practice occurred.

### **Rationale**

The framework used to guide the intervention is The Four-Level Model of the Healthcare System adapted from Ferlie and Shortell (2001). The conceptual model is shown in Appendix B. This model was chosen because it illustrates the concept of patient-centered care and its importance within the healthcare system. This model consists of four interconnected levels including the *Patient* at the center, surrounded by the *Care Team*, surrounded by the *Organization*, and finally encircled by the *Environment* (National Academy of Engineering and Institute of Medicine, 2005). It is crucial to identify the individual patient with their unique beliefs, desires, and needs as the nucleus for all healthcare interactions because “the role of the patient has changed from a passive recipient of care to a more active participant in care delivery” (National Academy of Engineering and Institute of Medicine, 2005). This framework calls upon healthcare providers to view patients and their families as “partners” allowing them to incorporate their needs and preferences into a free exchange of information with the healthcare team in which patients can communicate needs, participate in decision making, and assist in coordination of care (National Academy of Engineering and Institute of Medicine, 2005).

The Four-Level Model of the Healthcare System highlights the importance of patients and their support systems having access to tools, information, and education to help them navigate and integrate the critical but often overwhelming world of healthcare. This is the area in which the Care Team (anyone involved in delivering care to the patient) will implement the proposed intervention. The framework urges providers to be more responsive to the needs and preferences of patients which will be enhanced through the educational intervention developed in order to resolve the gap in present care.

A Plan-Do-Study-Act (PDSA) cycle was used to develop and implement the intervention. This concept comes from the Institute for Healthcare Improvement Model for Improvements and helps to focus and accelerate quality improvement initiatives (Agency for Healthcare Research and Quality, 2013). Planning this intervention included performing a literature review to identify gaps in practice and creating the idea for the project. Doing this intervention included carrying out the necessary steps to complete it. Studying this project included analyzing the data and results to determine if the aims were met. Acting after completion of this project included determining what worked well and what could be improved in future cycles. This intervention was expected to work because the concept of a continuous PDSA cycle allows researchers to make changes after results have been analyzed to improve the quality improvement initiative in the future. One PDSA cycle was utilized for the purpose of this project.

### **Purpose**

The purpose of this study was to assess the readability of local discharge information, present providers with practical tools to improve the readability of written instructions, and determine if a change of practice occurred. This was accomplished by utilizing the data collected in the PRIDE study (Balzer et al., 2019) to determine if there was a difference in patient

perception of the relative importance of discharge information, assessing the literacy level of the discharge information, and conducting an educational intervention for providers responsible for creating DIFs. The providers completed the Continuing Professional Development (CPD)-Reaction Questionnaire before and after the educational intervention to assess the impact of the intervention on clinical behavioral intentions (Légare et al., 2017). One-month post intervention a survey was administered to Advanced Practice Providers (APPs) to assess self-reported behavior change, and DIFs were assessed to determine if readability improved.

The aims were to:

- 1) Assess the literacy level of a random sample of DIFs used at patient discharge before and after the intervention.
- 2) Educate Advanced Practice Providers regarding the importance of literacy level and perceived importance of the information provided on the DIF.
- 3) Assess the impact of the intervention on clinical behavioral intentions and self-reported behavior change.

## **Methods**

### **Location**

The large urban academic medical center in which the intervention took place is a Joint Commission accredited level I trauma center containing over 800 beds (American Hospital Directory, 2018), and serving a vastly diverse patient population. This institution's mission statement is driven by quality and safety “to preserve and restore health for all people, to seek the cause and cure of diseases through innovative research, and to educate those who serve humanity”, with a vision “committed to excellence in patient care and education as the preeminent academic medical center in the mid-Atlantic region (Redacted, 2019). Over the last



three years, the medical center has discharged an average of 36,956 patients each year (Redacted, 2018).

### **Contextual Elements**

Contextual elements considered important at the outset of the intervention were communication between educator and recipient as well as learner willingness to change practice, because evidence-based practice relies both on disseminating the information and learner readiness to incorporate this information into practice (Young & Newell 2008). Other important contextual elements include recipient perception of educator's leadership, recipient perception of Doctor of Nursing Practice programs, and recipient patient/task load during an educational intervention. Understanding and managing these contextual elements is crucial because implementation research demonstrates that "when a majority of contextual elements are conducive to change, implementation is usually successful" (McCullough et al., 2015).

### **Ethical considerations**

This study was granted approval by both the Institutional Review Boards of the health system and James Madison University. This study posed no risks greater than those of everyday life to those involved and participation was entirely voluntary. Participation could be terminated at any time without penalty. All information collected was kept strictly confidential and was stored in a secure manner. Participants were notified of these facts in a written statement present on the questionnaires. Paper survey data were immediately entered into an excel spreadsheet maintained on a password protected computer within the institutional firewall of the health system. Paper surveys were shredded once information was entered into the spreadsheet. When this information needed to be transported off of the health system premises, an encrypted flash drive was used. The researcher reported no potential conflicts of interest.

## **Baseline Measures**

The initial assessment consisted of screening a random sample of fifty DIFs for grade level and readability. The random sample of DIFs was generated by the researcher monitoring the hospital discharge tracking system for patients flagged for discharge each day. Once a patient was flagged for discharge, the researcher screened the DIF to determine if it was created by a Hospital Medicine Advanced Practice Provider. Only DIFs created by Hospital Medicine APPs were utilized for inclusion.

After being identified for inclusion within the study, each DIF was assessed for grade level and readability using the Flesch-Kincaid Tool. The Flesch-Kincaid tool is the most widely used readability tool in the United States and provides users with two separate assessments (Flesch-Kincaid, 2019). The first assessment represents reading ease, and text is assigned a number between zero and one hundred using the formula  $206.835 - 1.015 \times (\text{words/sentences}) - 84.6 \times (\text{syllables/words})$  (Flesch-Kincaid, 2019). Low numbers represent complexity, therefore the higher the number, the lower the content complexity (Flesch-Kincaid, 2019). The second assessment provides users with the American school grade level required to understand the provided text using the formula  $0.39 \times (\text{words/sentences}) + 11.8 \times (\text{syllables/words}) - 15.59$  (Flesch-Kincaid, 2019). This data provided a baseline for comparison.

## **Intervention**

Next, an educational intervention was created and administered as an in-person poster presentation to ten Hospital Medicine APPs at an APP staff meeting. The APPs were identified through convenience sampling and included both Nurse Practitioners and Physician Assistants. The APPs were the recipients of the educational intervention because they are responsible for creating a majority of DIFs. A poster presentation format was chosen in order to summarize

information concisely and attractively, to help publicize information, and to generate discussion (NYU Libraries, 2018). This format increased the likelihood of reaching all department APPs because it was directly administered by the project director who was then available to answer any questions and participate in dialogue. Poster presentation dialogue has been shown to generate new ideas about applying or extending existing work, and raising new questions or suggestions to educate both the participants, and the researcher (Miller, 2007). This information helped to improve the intervention for future PDSA cycles. Prior to the staff meeting in which the presentation took place, an announcement email was sent to the APPs which can be seen in Appendix C.

The content of the poster presentation consisted of five sections including Abstract, Background, Our DIF Scores – March 2019, Action Steps, and How Will We Know if we are Successful. The Abstract section contained general information to introduce the project. The Background section discussed that most healthcare information is written at a reading level much too high for patients to effectively comprehend (Choudhry et al., 2016), the results of the PRIDE study (Balzer et al., 2019), and local health system information including rates of hospital discharges and readmissions. The Our DIF Scores – March 2019 section explained both the American Grade Level and Readability Score concepts and goal values, as well as contained graphs visually depicting the mean American Grade Level and Readability Scores of the surveyed DIFs. The Action Steps section contained practical suggestions for methods to improve grade level and readability, such as including the most important information first, limiting the number of messages, choosing words carefully, limiting jargon and technical terms, wording things in the positive, and emphasizing importance with bold (US Department for Health and Human Services & Centers for Disease Control and Prevention, 2009). The How Will We Know

if we are Successful section, cited improvements of follow up grade level and readability scores as well as provider self-report as measures of success. This structure allowed for future PDSA cycles to utilize or adapt the existing poster presentation to continue improving the process. The poster presentation can be seen in Appendix D.

### **Team**

The team involved in the work included a Doctor of Nursing Practice (DNP) student project director providing the educational intervention and assessing response, as well as the advanced practice providers employed within the Hospital Medicine service who were the recipients of the intervention. The team also included a faculty advisor from James Madison University who acted as Committee Chair, a statistician, and a terminal degree advisor from within the health system.

### **Outcome Measures**

The measures chosen include the CPD-Reaction questionnaire (Légare et al., 2017), the Flesch-Kincaid Readability Test Tool (Flesch-Kincaid, 2019), and a provider self-report survey (Légare et al., 2017). These measures helped meet the aims of assessing the average DIF literacy level, assessing the impact of the intervention, and assessing the effectiveness of the intervention. The rationale for choosing these measures was to reproduce a previously completed study. The operational definition of practice change was measured by a survey, and the operational definition of behavioral intention was measured by the CPD-Reaction tool. There were no specific measures to establish whether the observed outcomes were due to the intervention.

#### **CPD-Reaction Questionnaire**

First, the mean rankings of each CPD-Reaction construct were compared from before and after the intervention. The CPD-Reaction constructs are: *behavioral intention*, *social influence*,

*beliefs about capabilities, moral norm, and beliefs about consequences* (Légare et al., 2017). The construct *behavioral intention* assesses participants planning and intention to incorporate behavior change (Légare et al., 2017). The construct *social influence* measures participants beliefs surrounding whether peers do exhibit, or will exhibit, the targeted behavior (Légare et al., 2017). The construct *beliefs about capabilities* gauges participants confidence, in ease and ability to adopt behavior change (Légare et al., 2017). The construct *moral norm* evaluates how ethical and acceptable participants believe behavior change would be (Légare et al., 2017). The construct *beliefs about consequences* measures how useful and beneficial participants believe behavior change would be (Légare et al., 2017).

The CPD-Reaction Questionnaire is a valid and reliable “12-item instrument based on an integrated model combining a number of social cognitive theories for explaining health professionals’ clinical behavior through the proxy of intention” with Cronbach’s coefficients for the constructs varying from 0.79 to 0.89, as well as moderate test-retest reliability with weighted kappa values between 0.40 and 0.60 (Légare, Borduas, Freitas, & Turcotte, 2015). It also demonstrates construct validity with an exploratory factorial analysis confirming the presence of five constructs and a proportion of variance explained by each factor being superior to 5% (Légare et al., 2015).

This instrument is based on the assumption that three categories of variables predict health professionals’ behavior, 1) their intention to adopt a particular behavior or not; 2) their beliefs about their capabilities; and 3) their past behavior and habits (Légare et al., 2017). This tool has been successfully utilized in numerous populations including acute care nurses after a workplace violence training program (Lamont & Brunero, 2018), primary care providers after an online dementia care educational series (Bentley, Kerr, Ginger, & Karagoz, 2019), and acute care

providers after an educational intervention on non-pharmacological pain management (Booth, 2019). This tool assessed the impact of the educational intervention on clinical behavioral intentions, estimated the predictive potential for subsequent behavior change, and collected basic demographic information about participants.

### **Provider Self-Report**

Second, providers self-reported behavior changes as well as the percentage of clinical cases in which the behavior was adopted. This was used to assess the effectiveness of the intervention as providers were questioned regarding their perception on both improving the grade level and readability of discharge information, and incorporating patient preferences within discharge information. These questions were developed by the researcher and therefore there was no specific validity or reliability data regarding this tool.

### **Post-Intervention DIF Scores**

Third, DIFs before and after the intervention were compared for readability and grade level utilizing the Flesch-Kincaid Readability Tool (2019). This assessed for pre-intervention grade level and readability of DIFs, post-intervention grade level and readability of DIFs, to better understand the effectiveness of the intervention.

### **Data Collection Methods**

Immediately before and after the educational intervention, APPs completed a pen and paper survey containing the CPD-Reaction Questionnaire (Légare et al., 2017), which was adapted for this quality improvement project and can be found in Appendix E. Ten APPs were recruited to participate via convenience sampling by the researcher at an APP staff meeting.

One month after the intervention, those who initially completed the survey were contacted in person to complete a second pencil and paper survey to self-report behavior change

and estimate the percentage of clinical cases in which the behavior was adopted utilizing a four question Likert scale survey developed by the researcher. This survey can be found in Appendix F. At this time, a second sample of 50 DIFs was collected and evaluated utilizing the same procedure as the pre-intervention DIF assessment to determine if a change in practice occurred.

Informal ongoing assessment of the contextual elements of communication and learner willingness to change were monitored by the DNP student completing the educational intervention and administering the surveys. The student was available in person and by phone for questions or clarifications within the one month between the intervention and the follow up survey. No official notes or data were kept regarding this ongoing assessment.

The methods employed for ensuring completeness and accuracy of data collection included two individuals reviewing and recording the survey data, as well as an in-person review of surveys as they were completed to ensure all questions were answered. The methods employed for ensuring completeness and accuracy of data analysis include the researcher consistently utilizing the same Flesch-Kincaid website application for every DIF analysis.

### **Analysis**

SPSS version 24 (IBM Corporation, 2016) was used for data analysis. It was understood that variation within the data could occur with time due to both random variation and time trend variation. The random variation could not be controlled, however in an effort to control for time trend variation, follow up survey data as well as the second sample of DIFs were collected within a 6-day timeframe one month after the intervention.

### **CPD-Reaction Questionnaire**

Table 1 summarizes how items from the CPD-Reaction questionnaire were calculated according to the method created by Légare, and combined for analysis (Légare et al., 2017).

“Each item response format was pre-coded with Likert-type scale values (Table 1). The item score for each participant ranges from 1 to 7. A score for each construct was obtained by calculating the mean score for the construct (e.g., if the construct includes 2 items, the item scores were summed and divided by 2, yielding a score between 1 and 7)” (Légare et al., 2017).

Utilizing this formula, the questions “I intend to [behavior]” and “I plan to [behavior]” create the *behavioral intention* construct. The questions, “To the best of my knowledge, the percentage of my colleagues who [behavior] is”, “Now think about a co-worker whom you respect as a professional, in your opinion, will he/she [behavior]”, and “Most people who are important to me in my profession will [behavior]”, create the *social influence* construct. The questions “I am confident that I could [behavior] if I wanted to”, “For me, [behavior] would be”, and “I have the ability to [behavior]”, make up the *beliefs about capabilities* construct. The questions “[behavior] is the ethical thing to do”, and “It is acceptable to [behavior]”, make up the *moral norm* construct. Lastly, the questions “Overall, I think that for me [behavior] would be” useless to useful and harmful to beneficial, create the *beliefs about consequences* construct. The mean rankings of each of these constructs were compared before and after the intervention using a paired sample t-test.

Table 1.

*Construct Calculations*

Construct Scale		Items	Response Choices	Score by Construct
Intention	I <sub>1</sub>	I intend to [behavior]	Strongly disagree/agree	(I <sub>1</sub> +I <sub>7</sub> )/2
	I <sub>7</sub>	I plan to [behavior]	Strongly disagree/agree	
Social Influence	I <sub>2</sub>	To the best of my knowledge, the percentage of my colleagues who [behavior] is...	0-20%	(I <sub>2</sub> +I <sub>6</sub> +I <sub>9</sub> )/3
			21-40%	
			41-60%	
			61-80%	



			81-100%	
	I <sub>6</sub>	Now think about a coworker whom you respect as a professional. In your opinion, does he/she [ <i>behavior</i> ]?	Never/Always	
	I <sub>9</sub>	Most people who are important to me in my profession [ <i>behavior</i> ]	Strongly disagree/agree	
Beliefs about capabilities	I <sub>3</sub>	I am confident that I could [ <i>behavior</i> ] if I wanted to	Strongly disagree/agree	$(I_3+I_5+I_{11})/3$
	I <sub>5</sub>	For me, [ <i>behavior</i> ] would be...	Extremely difficult/easy	
	I <sub>11</sub>	I have the ability to [ <i>behavior</i> ]	Strongly disagree/agree	
Moral norm	I <sub>4</sub>	[ <i>Behavior</i> ] is the ethical thing to do	Strongly disagree/agree	$(I_4+I_{10})/2$
	I <sub>10</sub>	It is acceptable to [ <i>behavior</i> ]	Strongly disagree/agree	
Beliefs about consequences	I <sub>8</sub>	Overall, I think that for me [ <i>behavior</i> ] would be...	Useless/Useful	$(I_8+I_{12})/2$
	I <sub>12</sub>	Overall, I think that for me [ <i>behavior</i> ] would be...	Harmful/Beneficial	

Descriptive statistics such as frequencies, means with standard deviations, or medians with interquartile range were used to summarize participant characteristics and CPD-Reaction questionnaire item responses (Légare et al., 2017) for all quantitative data. Testing of normality was performed on the CPD-Reaction questionnaire responses including skewness and Kurtosis which demonstrated a slight deviation from the normal distribution. Skewness and Kurtosis measures were 1.36 and 0.11 for the *behavioral intention* construct, 0.92 and 3.10 for the *social influence* construct, 0.78 and -0.49 for the *beliefs about capabilities* construct, 0.89 and -0.16 for the *moral norm* construct, and 1.04 and -1.22 for the *beliefs about consequences* construct. Given that data were only slightly non-normally distributed, a paired sample t-test was used to compare the mean ranks of each construct obtained before and after professional development activities (Légare et al., 2017). To examine if the assumption of normal distribution might have affected our results, the Wilcoxon signed ranks test was also completed.

### Provider Self-Report

Nonparametric testing utilizing a Mann-Whitney test was performed to analyze DIF scores pre and post-intervention as these data were non-normally distributed. This test was utilized to interpret mean ranking. Data were treated as independent because one hundred different DIFs were sampled during a before and after time period.

### **Post-Intervention DIF Scores**

The self-reported behavior change surveys were reported as percentages as this information was only collected from participants at one point in time.

## **Results**

### **Demographic Information**

Overall, 10 participants completed the CPD-Reaction Questionnaire before and after the intervention with a 100% survey response rate (Table 1). 90% of respondents were female, 80% of respondents were Caucasian, with 60% of participants ranging in age from 25 to 34 years old. A majority of participants had practiced both as registered nurses for 1 to 5 years (70%), and as nurse practitioners for 1 to 5 years (80%).

Table 1. <i>Demographic Information</i>		
Characteristics	Category	N %
		Total = 10
Age (years)	25 - 34	6 (60)
	35 - 45	3 (30)
	45 - 54	0 ( 0)
	55 - 64	1 (10)

	65 - 74	0 ( 0)
Gender	Male	1 (10)
	Female	9 (90)
	Other	0 ( 0)
Race	Caucasian	8 (80)
	African American	2 (20)
	Hispanic	0 ( 0)
	Other	0 ( 0)
Years as RN	1 - 5	7 (70)
	6 - 10	2 (20)
	11 - 15	0 ( 0)
	16 - 20	0 ( 0)
	21 - +	1 (10)
Years as NP	1 - 5	8 (80)
	6 - 10	1 (10)
	11 - 15	1 (10)
	16 - 20	0 ( 0)
	21 - +	0 ( 0)

### CPD-Reaction Questionnaire

Each construct was compared before and after the educational intervention to determine if a change occurred. Mean results descriptively increased in all categories including *behavioral intention* (6.25 to 6.50), *social influence* (4.83 to 5.43), *beliefs about capabilities* (5.40 to 6.10), *moral norm* (6.35 to 6.65), and *beliefs about consequences* (6.45 to 6.60) (Table 3). Statistically

significant increases were found only in the area of *social influence* ( $p=0.040$ ) (Table 4). As the statistical results were similar between paired t-test and Wilcoxon signed rank test, results from the paired sample t-test are reported.

Table 3.

*Construct Comparison*

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	IntentionPre	6.25	10	1.03	0.33
	IntentionPost	6.50	10	0.85	0.27
Pair 2	SocialInfluencePre	4.83	10	1.06	0.33
	SocialInfluencePost	5.43	10	0.93	0.29
Pair 3	CapabilitiesPre	5.40	10	0.94	0.30
	CapabilitiesPost	6.10	10	0.74	0.23
Pair 4	MoralNormPre	6.35	10	0.78	0.25
	MoralNormPost	6.65	10	0.63	0.20
Pair 5	ConsequencesPre	6.45	10	0.80	0.25
	ConsequencesPost	6.60	10	0.70	0.22

Table 4.

*Paired Sample Test*

		Mean	Std. Dev.	Std. Error Mean	95% Confidence Interval		t	df	Sig. (2-tailed)
					Lower	Upper			
Pair 1	IntentPre IntentPost	-0.25	0.42	0.13	-0.55	0.05	-1.86	9	0.10
Pair 2	SocialPre SocialPost	-0.60	0.77	0.24	-1.15	-0.05	-2.48	9	<b>0.04</b>
Pair 3	CapablePre CapablePost	-0.70	1.12	0.35	-1.50	0.10	-1.98	9	0.08
Pair 4	MoralPre MoralPost	-0.30	0.71	0.23	-0.81	0.21	-1.33	9	0.22
Pair 5	ConsequencePre ConsequencePost	-0.15	0.24	0.08	-0.32	0.02	-1.96	9	0.08

### Provider Self-Report

Of the original 10 participants, 100% completed the one month follow up survey to self-report behavior change in the areas of improving DIF readability and incorporating patient preferences. 80% of APPs reported that they agreed or strongly agreed that they had increased the readability of their DIFs, and 40% felt that they had improved readability in 80-100% of clinical cases in the preceding 30 days. 70% of APPs reported that they agreed or strongly agreed that they had incorporated patient preferences into their DIFs, and 60% felt that they had incorporated patient preferences in 80-100% of clinical cases in the previous 30 days.

Question	Category	N	%
		Total = 10	
In the last month I have increased the readability of my DIFs:	Strongly Agree	4	(40)
	Agree	4	(40)
	Somewhat Agree	1	(10)
	Neither Agree nor Disagree	1	(10)
	Somewhat Disagree	0	( 0)
	Disagree	0	( 0)
	Strongly Disagree	0	( 0)
The percentage of clinical cases in which I improved the	81 - 100%	4	(40)
	61 - 80%	3	(30)
	41 - 60%	2	(20)
	21 - 40%	1	(10)

readability of my DIFs:	0 - 20%	0 ( 0)
In the last month I have incorporated patient preferences in my DIFs:	Strongly Agree	5 (50)
	Agree	2 (20)
	Somewhat Agree	1 (10)
	Neither Agree nor Disagree	2 (20)
	Somewhat Disagree	0 ( 0)
	Disagree	0 ( 0)
	Strongly Disagree	0 ( 0)
The percentage of clinical cases in which I incorporated patient preferences in my DIFs:	81 - 100%	6 (60)
	61 - 80%	1 (10)
	41 - 60%	2 (20)
	21 - 40%	1 (10)
	0 - 20%	0 ( 0)

### Pre/Post-Intervention DIF Scores

Initial DIF assessments demonstrated a mean grade level of 9.5 ranging from 8<sup>th</sup> to 16<sup>th</sup> grade. Initial readability scores demonstrated a mean of 61.70 ranging from 35.80 to 67.70. Analysis of 50 DIFs after the educational intervention, found that mean grade level decreased from 9.5 to 7.2 with range decreasing from 8<sup>th</sup> to 16<sup>th</sup> grade to a range of 5<sup>th</sup> to 10<sup>th</sup> grade. Mean readability scores increased from 61.70 to 64.25 with a range increasing from 35.80 to 67.70 to a range of 55.90 to 74.40. Statistical differences are explained with the Mann-Whitney Ranks (Table 5), and Mann-Whitney Testing (Table 6). Statistically significant changes were found in grade level ( $p=0.001$ ), and near statistically significant changes were found in readability ( $p=0.051$ ).

Table 5.

*Mann-Whitney Ranks*

	<b>Time</b>	<b>N</b>	<b>Mean Rank</b>	<b>Sum of Ranks</b>
DIFs	Pre	50	50.50	2525.00
	Post	50	50.50	2525.00
	Total	100		
Grade Level	Pre	50	60.32	3016.00
	Post	50	40.68	2034.00
	Total	100		
Readability	Pre	50	44.84	2242.00
	Post	50	56.16	2808.00
	Total	100		

Table 6.

*Mann-Whitney Testing*

	<b>DIFs</b>	<b>Grade Level</b>	<b>Readability</b>
Mann-Whitney U	1250.00	759.00	967.00
Wilcoxon W	2525.00	2034.00	2242.00
Z	0.00	-3.39	-1.10
Asymp. Sig. (2-tailed)	1.00	<b>0.001</b>	<b>0.051</b>

**Discussion**

The Joint Commission has set literacy benchmarks for hospitals to achieve (The Joint Commission, 2007), the Institute of Medicine has set improving health literacy as a crucial quality improvement goal (Institute of Medicine, 2004), and other highly influential organizations such as the American Medical Association (1999) have recommended enhanced research surrounding health literacy and increased education for the medical community since as early as 1999. With these recommendations and the increased focus of health systems on preventing readmissions, hospitals and providers are motivated to incorporate new ideas that could generate sustainable change in these areas of healthcare. Despite these recommendations,

literature review demonstrated a lack of training for providers regarding strategies to meet these benchmarks.

At baseline review of the hospital discharge process within a large academic health system, it was noted that although a discharge template was standardized, great variability took place within the grade level and readability scores of the discharge information being given to patients. This supports findings in the literature which demonstrate a wide range of grade-level and readability scores within discharge information (Choudhry et al., 2016), as well as an overall inconsistency within the discharge process (Buikstra, Strivens, & Clay-Williams, 2020).

Patients have the right to consistently receive healthcare information in a way they can understand, and providers have the right to receive education on how to provide this service to their patient populations. By prioritizing patient preferences and incorporating simple strategies, providers can make impactful changes to optimize patient comprehension and learning. This study investigated the effect of an educational intervention on hospital discharge information grade level and readability as well as the behavioral intentions of those writing the discharge information. This study led to important findings within the areas of the CPD-Reaction Questionnaire, the Provider Self-Report Survey, and the Post-Intervention DIF Scores.

### **CPD-Reaction Questionnaire**

The one construct within the CPD-Reaction Questionnaire (Légare et al., 2017) which demonstrated a statistically significant change, the *social influence* construct, consisted of three questions. These questions were “To the best of my knowledge, the percentage of my colleagues who will improve the readability of their DIFs is”, “Now think about a co-worker whom you respect as a professional, in your opinion, will he/she improve the readability of their DIFs”, and “Most people who are important to me in my profession will improve the readability of their



DIFs”. The change in this area after the intervention demonstrates a strong belief among APPs that colleagues will incorporate a behavior change.

These findings support existing evidence because literature has demonstrated that “nurses, particularly advanced practice nurses, are socialized into a hierarchy that has implicit values and roles” (Waugaman & Lohrer, 2000) in which emulating and adopting the behaviors of respected peers is common (Felstead & Springett, 2016). If key leaders within the nursing hierarchy adopt a practice, it is much more likely for peers to incorporate these behaviors and sustain practice change. This is especially important when considering the demographics of this particular study group, which consisted primarily of advanced practice nurses very early in their careers. Transitioning to the advanced practice role can be a tumultuous time, and nurses often idealize experienced nursing mentors and seek to emulate their practice (Ferguson, 2011). By generating the belief that peers will change practice, this intervention becomes more likely to create sustainable change, and adds to the existing literature by demonstrating the importance of providers receiving this education early within their advanced practice role.

Although the only construct to demonstrate significant change was *social influence*, the remaining constructs of *behavioral intention*, *beliefs about capabilities*, *moral norm*, and *beliefs about consequences* each showed a mean increase post-intervention. An increase in the *behavioral intention* construct supports what is known, in that education can increase the likelihood of intending to adopt behavior change (Saddawi-Konefka et al., 2016). However, even a statistically significant change to this construct would be unlikely to generate true practice change, given the often-large gap between intention and action (Saddawi-Konefka et al., 2016). An increase in the *beliefs about capabilities* construct supports existing evidence, which demonstrates that receiving education about incorporating a behavior increases confidence in the

ability to incorporate the behavior (Terry & Cutter, 2015). An increase in the *moral norm* construct also supports existing literature, in that increasing the belief that an action is morally and ethically acceptable, makes it more likely to be adopted, especially within a medical community in which moral and ethical standards are deeply engrained into education (Haddad & Geiger, 2019). An increase in the *beliefs about consequences* construct supports the concept of incorporating evidence-based practice, in that providers are more likely to integrate current best evidence when they believe a behavior will be beneficial and helpful (Titler, 2008).

### **Provider Self-Report**

No previous studies have been conducted in which providers self-report behavior changes in the domains of incorporating patient preferences and improving the readability of discharge information after an educational intervention, therefore these results add to what is known. Providers self-reported a perceived improvement in both incorporating patient preferences and improving the readability of their discharge information forms in a majority of clinical cases one month after the intervention. Given that actual change was demonstrated in the post-intervention DIF scores, this study suggests that measuring self-report behavior changes may be helpful for understanding subsequent behavior.

### **Post-Intervention DIF Scores**

National benchmarks state that information provided to patients should be written at a 6th grade level or below (Choudhry et al., 2016), however many available patient education materials are written at a level that not only does not meet these standards, but are likely too complex for a substantial portion of the population to comprehend (Badarudeen & Sabharwal, 2010). As seen on the initial DIF assessment, the Hospital Medicine Service within the health system in which the intervention took place, was also far above meeting these benchmarks which

supports existing literature. Initial data demonstrated a mean DIF grade level of 9.5, which was 3.5 grade levels over the recommended levels. After the intervention, mean grade level dropped to 7.2, only 1.2 grade levels over the goal. Although the mean grade level scores did not meet the national recommendations of 6th grade or below, the post-test levels did surpass assessments completed by others, which frequently demonstrated discharge information written at college graduate reading levels or above (Choudhry et al., 2016). With repeated PDSA cycles and continuous provider education, it may be possible to meet national benchmarks within the Division of Hospital Medicine. Further research needs to be completed to determine if this education would be successful in other disciplines.

Many disciplines argue that the information that must be conveyed to patients is too complex to summarize within the national benchmarks, and that oversimplifying text can make the material too straightforward to convey necessary information (Badarudeen & Sabharwal, 2010). In contrast to this viewpoint, several studies have demonstrated that improving the readability of patient information leads to improved patient outcomes (Badarudeen & Sabharwal, 2010). The findings of this study support that the 6<sup>th</sup> grade benchmark may be unattainable with current methods and resources. The question of whether patients fully comprehend the key facts within text written at a 6<sup>th</sup> grade level needs to be explored. Other aspects that need to be explored within the institution include whether DIFs written at a 6<sup>th</sup> grade level improve post-hospital follow up adherence, as was demonstrated by Marcentoni, et al. (2015), and whether a benefit could be found by creating disease-specific instructions templates written at a 6<sup>th</sup> grade level (Mueller et al., 2015).

In the local health system in which the intervention took place, some social workers screen patients for the highest educational grade level attained in their admission assessments.

However, the problem with this approach is that patients typically demonstrate reading levels of five full grades lower than their highest attained educational grade (Badarudeen & Sabharwal, 2010). The average resident of the United States reads at an 8<sup>th</sup> grade reading level, and the average Medicare beneficiary reads at a 5<sup>th</sup> grade reading level (Stossel, Segar, Gliatto, Fallar, & Karani, 2012). Given these facts, and the general mismatch between patient reading skills and the readability of health information, some interventions aimed at improving written health information have focused on screening individual patients for reading levels using tools such as Rapid Estimate of Adult Literacy in Medicine (REALM), Test of Functional Health Literacy in Adults (TOFHLA), or Newest Vital Sign (NVS) (Badarudeen & Sabharwal, 2010). Further research needs to be done on creating a combined approach consisting of assessing literacy level and delivering tailored materials based on this assessment in order to facilitate truly patient-centered care.

While the grade level and readability scores did improve significantly, there are still substantial areas for improvement within the discharge process. For example, providers are attempting to incorporate patient preferences by placing medication information first within the discharge information section. However, when the actual discharge packet is printed by nurses, the discharge information section is imbedded within numerous pages of auto populated text containing information that is redundant and not applicable to many patients. Furthermore, strategies demonstrated to improve understanding of discharge instructions and patient satisfaction such as the teach-back method (Burke, 2018), “do not disturb signs” (Polster, 2015), and a patient navigator (Kwan et al., 2013), are not utilized consistently by nursing staff within the institution. Although this project focused on improving the readability of discharge

information content, health systems should consider multiple strategies to improve all aspects of the discharge process.

### **Limitations**

The main limitation of this study is the small sample size of participants. A power analysis demonstrated that for future PDSA cycles, at least 34 participants would be ideal. However, at the time of this study, there were only 16 Hospital Medicine APPs employed by the health system, and only 10 were available for study participation. It also needs to be considered that despite the small sample size, a statistically significant change was achieved, meaning that more outcomes could have achieved statistical significance with a larger sample size.

Another limitation of this study is that the educational intervention was only administered to 10 out of the 16 Hospital Medicine APPs. This means that the post-intervention DIFs surveyed were created by a combination of both APPs who did and did not receive the education. Therefore, the total effect size was diluted, and the true effect size was likely even more substantial than what was represented within the data.

Another limitation is the use of one tool to check both grade level and readability of discharge information. Numerous applications utilize different mathematical equations to calculate these factors. Criticism is drawn because these tools work under the assumption that longer words and sentences equate to increasing complexity, the fact that the tools are unable to assess the active role of the reader, and the fact that different formulas can generate vastly different results for the same text (Agency for Healthcare Research and Quality, 2015). Given these factors, the Flesh-Kincaid tool was chosen because it is widely regarded as the most popular and accurate of these applications.

### **Conclusion**

The APP education proved to be a useful intervention in improving discharge communication grade level and readability, substantially contributing to existing literature. A significant change was made to APP social influence beliefs, their self-reported behavior change, and DIF grade level and readability scores. Continued education and PDSA cycles are needed to ensure sustainability of the demonstrated practice change in an environment that can be subject to high staff turnover. Further research needs to be done utilizing alternative patient populations and larger sample sizes to determine if improved readability correlates to decreased hospital readmissions.

Providers responsible for creating patient discharge information should be equipped with the knowledge of how to write literacy level appropriate materials. It is the responsibility of the health system to provide this education. Considering the results of this quality improvement project, there is an opportunity to improve discharge information readability through inpatient provider education.

## Appendix A

### Patient Perceptions of Relative Importance of Discharge Elements (PRIDE) Study

We are requesting your participation in a research survey to understand how you value various components of the discharge information (or paperwork) you receive. This survey will help us to determine which component of the discharge information is most important to our patients. The survey will take approximately 3 to 5 minutes of your time. We anticipate no risks to you if you participate in this study. Your participation in this survey is completely voluntary. If you choose to take the survey, but do not wish to complete it, you can stop at any time. Your decision to participate in this survey will not affect your care. All collected information will be kept strictly confidential and will be stored in a secure manner. We will not share any individual's data and only summary results from this study will be used for publication.

There is no compensation for the completion of this survey. If you have any questions, you can contact the primary investigator, Dr. Rehan Qayyum at 804-628-3624. You must be at least 18 years of age or older to participate in this study. By beginning this survey you indicate your consent for participation.

We thank you for your participation!

**Directions:** There are 7 categories listed below, one of which is listed as "other". Please rank the categories from # 1 to # 7 (with #1 being the most important and # 7 being the least important to you). Each number from 1 to 7 should only be used once. You can use the "other" category to include things that you feel should be included, but wasn't mentioned here.

**Medications** Rank: \_\_\_\_\_  
*(What medications to take and how to take them?)*

**Summary of Hospital Stay** Rank: \_\_\_\_\_  
*(Why you were hospitalized and what was done during your hospital stay?)*

**Specific Instructions for Your Disease** Rank: \_\_\_\_\_  
*(Such as checking your blood pressure, blood glucose, weight, etc.)*

**Activity/Diet Restrictions** Rank: \_\_\_\_\_  
*(What physical activity to do or what foods to avoid?)*

**When to call 911/** **Doctor Rank:** \_\_\_\_\_  
*(Symptoms that should prompt a call to 911 or your primary doctor)*

**Follow-up** **Appointments Rank:** \_\_\_\_\_  
*(When, where, and who you will be seeing after discharge?)*

**Other** \_\_\_\_\_ **Rank:** \_\_\_\_\_

**Age:** \_\_\_\_\_ *(Please use 89 if you are older than 89 years)*

**Race (circle one):**

African American      Caucasian      Hispanic      Other

**Gender (circle one):**

Male                      Female

**Education Level (circle one):** None      Elementary School      Middle School      High School

College      Post-graduate

**Number of Home Medications:** \_\_\_\_\_

**Number of Medication Changes at Discharge:** \_\_\_\_\_

**Discharge Disposition (circle one):**

Home      Nursing Facility      Hospice      Rehabilitation

Long term acute care center                      Other



**Medical Conditions (circle all that apply):**

Diabetes Mellitus

Hypertension

Heart Failure

Chronic Kidney Disease

COPD

Asthma

Tobacco Use

List any other conditions here:


---

---

---

---

**Appendix B**



(Ferlie & Shortell, 2001)

### **Appendix C**

Dear Hospital Medicine Advanced Practice Providers,

At the end of the upcoming APP Staff Meeting I will be performing a brief educational intervention for the group to disseminate both the results of the PRIDE study and the results of our average DIF grade level scores. I will also share some strategies to make our DIFs more patient centered and readable. I will ask you to perform a brief survey before and after the educational session as part of a research project. Participation in this research is entirely voluntary.

Thank you in advance for your attendance and participation!

Appendix D


# Improving Readability of Discharge Instructions

Amber Balzer, MSN, APRN, CMSRN, AGACNP-BC  
Erica Lewis, RN, PhD



**Abstract**  
Patient instructions should be written at or below a 6th grade reading level. Our reading levels are too high. In order to change this, we need to change the way we word what we write in our Discharge Instruction Forms (DIFs). Practical suggestions are made for how we can improve. Small changes can make a big difference, keeping our patient's safe and helping them to avoid readmission. We will know we succeed if screening of DIFs reveals decreased grade level, increased readability scores, and if providers self-report behavior change post-intervention.

**Background**  
•According to the National Institute of Health, all patient instructions should be written at or below a 6th-grade level to help prevent readmission. Yet, nationwide most discharge instructions are written at college graduate reading level (Choudhry, et al, 2016).  
•In the PRIDE study, our patient's ranked medication instructions as the most important discharge element.  
•Despite being responsible for over 19% of total health system discharges, our division's current DIFs lack individualization and our DIF literacy level was previously unknown.  
•Our health system readmission rate is approaching 16%.



- Action Steps**
- Include the most important information first (CDC)
  - State actions to take  
Say why it's important
  - Limit the number of messages
  - Focus on what the patient needs to know  
Skip excess details  
Avoid lengthy lists  
Use bullets instead of commas
  - Choose words carefully
  - Use words with 1-2 syllables  
Use sentences with 8-10 words  
Paragraphs should be only 3-5 sentences
  - Limit jargon and technical terms
  - When possible, word things in the positive
- Saying what patients should do instead of saying what they should not do
- Use bold to emphasize importance, avoid all caps, italics, and underlines

- How will we know if we are successful?**
- Repeat screening of DIFs reveals decreased grade level
  - Repeat screening of DIFs reveals increased reading ease
  - Providers self-report behavior change

## Appendix E

I am requesting your participation in a research study to help determine your behavioral intentions surrounding the discharge process, and if a change in behavioral intention can take place. The research activities include:

- 1) The researcher will assess a random sample of DIFs for readability and grade level
- 2) Participants will complete a behavioral intention survey before an educational intervention
- 3) The researcher will present participants with an in person educational intervention focused on improving patient satisfaction with the DIF by incorporating prioritized patient preferences and making the DIF more readable by utilizing appropriate literacy levels
- 4) Participants will complete a behavioral intention survey after the educational intervention
- 5) The researcher will then assess a random sample of DIFs for readability and grade level
- 6) One month after the intervention the researcher will interview participants about practice Change

The surveys and interviews will take approximately 3 to 5 minutes of your time, and the educational intervention will take approximately 15 minutes of your time. I anticipate no risks to you if you participate in this study. Your participation in this study is completely voluntary. If you choose to take the survey, but do not wish to complete it, you can stop at any time. All collected information will be kept strictly confidential and will be stored in a secure manner. I will not share any individual's data and only summary results from this study will be used for publication.

There is no compensation for the completion of this study. If you have any questions, you can contact the primary investigator, Amber Balzer at 804-297-5294. By beginning this survey, you indicate your consent for participation.

Thank you for your participation!

**Directions:** Please answer each of the following questions by indicating the number that best describes your opinion about the behavior indicated. Some of the questions may appear to be similar, but they do address somewhat different aspects of the behavior stated.

1. I intend to improve the readability of my DIFs: Strongly disagree Strongly agree  
[1] [2] [3] [4] [5] [6] [7]

2. To the best of my knowledge, the percentage of my colleagues who will improve the readability of their DIFs is: 0-20% 21-40% 41-60% 61-80% 81-100%  
[1] [2] [3] [4] [5]

3. I am confident that I could improve Strongly disagree Strongly agree

the readability of my DIFs if I wanted to: [1] [2] [3] [4] [5] [6] [7]

---

4. Improving the readability of my DIFs is the ethical thing to do: Strongly disagree [1] [2] [3] [4] [5] [6] [7] Strongly agree

---

5. For me, improving the readability of my DIFs would be: Extremely difficult [1] [2] [3] [4] [5] [6] [7] Extremely easy

---

6. Now think about a co-worker whom you respect as a professional, in your opinion, will he/she improve the readability of their DIFs? Never [1] [2] [3] [4] [5] [6] [7] Always

---

7. I plan to improve the readability of my DIFs: Strongly disagree [1] [2] [3] [4] [5] [6] [7] Strongly agree

---

8. Overall, I think that for me improving the readability of my DIFs would be: Useless [1] [2] [3] [4] [5] [6] [7] Useful

---

9. Most people who are important to me in my profession will improve the readability of their DIFs: Strongly disagree [1] [2] [3] [4] [5] [6] [7] Strongly agree

---

10. It is acceptable to improve the readability of my DIFs: Strongly disagree [1] [2] [3] [4] [5] [6] [7] Strongly agree

---

11. I have the ability to improve the readability of my DIFs: Strongly disagree [1] [2] [3] [4] [5] [6] [7] Strongly agree

---

12. Overall, I think that for me improving the readability of my DIFs would be: Harmful [1] [2] [3] [4] [5] [6] [7] Beneficial

---

Demographics:

Age: [25-34] [35-44] [45-54] [55-64] [65-74]

Gender: [Male] [Female] [Other]

Race:      [Caucasian]   [African American]   [Hispanic]   [Other]

Years Practicing as Registered Nurse:      [1-5]   [6-10]   [11-15]   [16-20]   [21 or more]

Years Practicing as Nurse Practitioner:      [1-5]   [6-10]   [11-15]   [16-20]   [21 or more]

### Appendix F

Directions: Please answer each of the following questions by indicating the response that best describes your practice.

1. In the last month I have increased the readability of my DIFs:	Strongly disagree					Strongly agree	
	[1]	[2]	[3]	[4]	[5]	[6]	[7]

---

2. The percentage of clinical cases in which I improved the readability of my DIFs:	0-20%	21-40%	41-60%	61-80%	81-100%
	[1]	[2]	[3]	[4]	[5]

---

3. In the last month I have incorporated patient preferences in my DIFs:	Strongly disagree					Strongly agree	
	[1]	[2]	[3]	[4]	[5]	[6]	[7]

---

4. The percentage of clinical cases in which I incorporated patient preferences in my DIFs:	0-20%	21-40%	41-60%	61-80%	81-100%
	[1]	[2]	[3]	[4]	[5]

---



## References

- Agency for Healthcare Research and Quality. (2013). Plan-Do-Study-Act (PDSA) Cycle. Retrieved from <https://innovations.ahrq.gov/qualitytools/plan-do-study-act-pdsa-cycle>.
- Agency for Healthcare Research and Quality. (2018). Readmissions and Adverse Events After Discharge. Retrieved from <https://psnet.ahrq.gov/primers/primer/11/Readmissions-and-Adverse-Events-After-Discharge>.
- Agency for Healthcare Research and Quality. (2013). Re-Engineered Discharge (RED) Toolkit. Retrieved from <https://www.ahrq.gov/professionals/systems/hospital/red/toolkit/redtool4.html>.
- Agency for Healthcare Research and Quality. (2015). Use Caution with Readability Formulas for Quality Reports. Retrieved from <https://www.ahrq.gov/talkingquality/resources/writing/tip6.html>.
- Albrecht, J. S., Gruber-Baldini, A. L., Hirshon, J. M., Brown, C. H., Goldberg, R., Rosenberg, J. H., ... Furuno, J. P. (2014). Hospital Discharge Instructions: Comprehension and Compliance Among Older Adults. *Journal of General Internal Medicine*, 29(11), 1491-1498. Retrieved from <https://link.springer.com/article/10.1007/s11606-014-2956-0>.
- American Hospital Directory. (2018). VCU Medical Center. Retrieved from [https://www.ahd.com/free\\_profile/490032/VCU\\_Medical\\_Center/Richmond/Virginia/](https://www.ahd.com/free_profile/490032/VCU_Medical_Center/Richmond/Virginia/).
- American Medical Association Ad Hoc Committee on Health Literacy for the Council on Scientific Affairs (1999). Health Literacy: Report of the Council on Scientific Affairs *JAMA*, 281(6), 552–557. doi:10.1001/jama.281.6.552.
- Balzer, A., Kurbanova, N., Reiber, C., Farooq, F., Uppalapati, S., Gwalani, P., Qayyum, R.

- (2019, March). *Patients' Perceptions of Relative Importance of Discharge Elements*. Poster session presented at the Society of Hospital Medicine Annual Conference, Alexandria, VA.
- Badarudeen, S., & Sabharwal, S. (2010). Assessing Readability of Patient Education Materials: Current Role in Orthopaedics. *Clinical Orthopedics and Related Research*, 468(10), 2572-2580. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3049622/>.
- Bentley, M. W., Kerr, R., Ginger, M., & Karagoz, J. (2019). Behavioural change in primary care professionals undertaking online education in dementia care in general practice. *Australian Journal of Primary Health*, 25(3), 244-249. Retrieved from <http://www.publish.csiro.au/py/PY18079>.
- Blaine, K., Rogers, J., O'Neil, M. R., McBride, S., Faerber, J., Feudtner, C., & Berry, J. G. (2018). Clinician Perceptions of the Importance of Hospital Discharge Components for Children. *Journal for Healthcare Quality*, 40(2), 79-88. Retrieved from [https://www.nursingcenter.com/journalarticle?Article\\_ID=4564811&Journal\\_ID=3304723&Issue\\_ID=4564750](https://www.nursingcenter.com/journalarticle?Article_ID=4564811&Journal_ID=3304723&Issue_ID=4564750).
- Booth, L. S. (2019). Provider Attitudes on Using Nonpharmacological Methods for Pain Management in Acute Care - Post an Educational Intervention. *The University of Arizona, ProQuest Dissertations*. Retrieved from <https://search.proquest.com/openview/a14d97b2c0f644ce65728cc584c90b6a/1?pq-origsite=gscholar&cbl=18750&diss=y>.
- Boston University. (2014). Project Red. Re-Engineered Discharge. Retrieved from <https://www.bu.edu/fammed/projectred/index.html>.
- Buikstra, B., Strivens, S., & Clay-Williams, R. (2020). Understanding variability in discharge

- planning processes for the older person. *Safety Science*, 121, 137-146. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0925753518320538>.
- Burke, K. (2018). Improving Patient Discharge Satisfaction Scores by Implementing Teach-Back Instructions in a Community Hospital Emergency Department (ED): A Quality Improvement Project. *Doctor of Nursing Practice (DNP) Projects*, 160. Retrieved from [https://scholarworks.umass.edu/nursing\\_dnp\\_capstone/160](https://scholarworks.umass.edu/nursing_dnp_capstone/160).
- Cawthon, C., Walia, S., Osborn, C. Y., Niesner, K. J., Schnipper, J. L., & Kripalani, S. (2013). Improving Care Transitions: The Patient Perspective. *Journal of Health Communication*, 17(3), 312-324. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3603351/>.
- Choudhry, A. J., Baghdadi, Y. M. K., Wagie, A. E., Habermann, E. B., Heller, S. F., Jenkins, D. H., ... Zielinski, M. D. (2015). Readability of Discharge Summaries: With What Level of Information are we Dismissing our Patients? *The American Journal of Surgery*, 211(3), 631-636. Retrieved from [https://www.researchgate.net/publication/288701294\\_Readability\\_of\\_Discharge\\_Summaries\\_With\\_What\\_Level\\_of\\_Information\\_are\\_we\\_Dismissing\\_our\\_Patients](https://www.researchgate.net/publication/288701294_Readability_of_Discharge_Summaries_With_What_Level_of_Information_are_we_Dismissing_our_Patients).
- Corser, W., Dontje, K., Neuberger, N., Chant, E., & Keskimaki, A. (2017). "After Visit Patient Summaries": A Patient Perspective. *Journal of Nursing Informatics*, 21. Retrieved from <https://www.himss.org/library/after-visit-patient-summaries-patient-Perspective>.
- El-Eid, G. R., Kaddoum, R., Tamim, H., & Hitti, E. A. (2015). Improving Hospital Discharge Time. *Medicine (Baltimore)*, 94(12), 633. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4554014/>.

- Felstead, I. S., & Springett, K. (2016). An exploration of role model influence on adult nursing students' professional development: A phenomenological research study. *Nurse Education Today*, 37, 66-70. Retrieved from <https://www.sciencedirect.com/science/article/abs/pii/S0260691715004785>.
- Ferguson, L. M. (2011). From the perspective of new nurses: What do effective mentors look like in practice? *Nurse Education in Practice*, 11(2), 119-123. Retrieved from <https://www.sciencedirect.com/science/article/abs/pii/S1471595310001502>.
- Ferlie, E. B., & Shortell, S. M. (2001). Improving the quality of health care in the United Kingdom and the United States: a framework for change. *Milbank Quarterly*, 79(2), 281–315. [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2751188/pdf/milq\\_206.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2751188/pdf/milq_206.pdf).
- Flesch-Kincaid. (2019). Readability Test Tool. Retrieved from <https://www.webfx.com/tools/read-able/flesch-kincaid.html>.
- Haddad, L. M., & Geiger, R. A. (2019). Nursing Ethical Considerations. *StatPearls*. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK526054/>.
- Health Literacy Basics. (2017). Quick Guide to Health Literacy. Retrieved from <https://health.gov/communication/literacy/quickguide/factsbasic.htm>.
- Horwitz, L. I., Jenq, G. Y., Brewster, U. C., Chen, C., Kanade, S., Van Ness, P. H., ..... Krumholz, H., M. (2013). Comprehensive quality of discharge summaries at an academic medical center. *Journal of hospital medicine*, 8(8), 436-43.
- IBM Corporation. (2016). IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp. Retrieved from <https://www01.ibm.com/support/docview.wss?uid=swg21476197>.
- Institute for Healthcare Improvement. (2019). Post Discharge Tool. Retrieved from <https://www.npsf.org/page/postdischargetool>.
- Institute of Medicine. (2004). Health Literacy: A Prescription to End Confusion. *Washington*,

- DC: *The National Academies Press*. Retrieved from <http://www.nationalacademies.org/hmd/Reports/2004/Health-Literacy-A-Prescription-to-End-Confusion.aspx>.
- Jakucs, C. (2018). Discharge planning starts at admission. *Nurse.com*. Retrieved from <https://resources.nurse.com/discharge-planning-starts-admission>.
- Kwan, J. L., Bell, C. M., Morgan, M. W., & Stewart, T. E. (2013). Inpatient Patient Navigator Program Reduces Length of Stay. *Harvard Business Review*. Retrieved from <https://hbr.org/2013/11/inpatient-patient-navigator-program-reduces-length-of-stay>.
- Lamont, S. & Brunero, S. (2018). The effect of a workplace violence training program for generalist nurses in the acute hospital setting: A quasi-experimental study. *Nurse Education Today*, 68, 45-52. Retrieved from <https://www.sciencedirect.com/science/article/abs/pii/S0260691718302004>.
- Légare, F., Borduas, F., Freitas, A., & Turcotte, S. (2015). The Continuing Professional Development (CPD) Reaction Questionnaire User Manual.
- Légare, F., Freitas, A., Turcotte, S., Borduas, F., Jacques, A., Luconi, F., ... Labrecque, M. (2017). Responsiveness of a simple tool for assessing change in behavioral intention after continuing professional development activities. *PLoS ONE*, 12(5). Retrieved from <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0176678>.
- Marcantoni, J. R., Finney, K., & Lane, M. A. (2015). Using Health Literacy Guidelines to Improve Discharge Education and the Post-Hospital Transition: A Quality Improvement Project. *American Journal of Medical Quality*, 29(1), 86. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3986412/>.
- McCullough, M. B., Chou, A. F., Solomon, J. L., Petrakis, B. A., Kim, B., Park, A. M.,

- Benedict, A. J., ...Rose, A. J. (2015). The interplay of contextual elements in implementation: an ethnographic case study. *BMC Health Serv Res*, 15(62). Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4345021/>.
- Miller, J. (2007). Preparing and Presenting Effective Research Posters. *Health Services Research*. 42(1), 311-328. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1955747/>.
- Mueller, S. K., Giannelli, K., Boxer, R., & Schnipper, J. L. (2015). Readability of patient discharge instructions with and without the use of electronically available disease-specific templates. *Journal of American Medical Association*, 22, 857-863. doi:10.1093/jamia/ocv005.
- National Academy of Engineering, Institute of Medicine, Committee on Engineering and the Health Care System. (2005). Building a Better Delivery System: A New Engineering/Health Care Partnership. *Washington (DC): National Academies Press*. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK22878/>.
- NEJM Catalyst. (2018). Hospital Readmissions Reduction Program (HRRP). *New England Journal of Medicine Catalyst*. Retrieved from <https://catalyst.nejm.org/hospital-readmissions-reduction-program-hrrp/>.
- NYU Libraries. (2018). How to Create a Research Poster: Poster Basics. Retrieved from <https://guides.nyu.edu/posters>.
- Polster, D. (2015). Preventing readmissions with discharge education. *Nursing Management*, 46(10), 30-37. Retrieved from [https://www.nursingcenter.com/cearticle?an=00006247-201510000-00007&Journal\\_ID=54013&Issue\\_ID=3208683](https://www.nursingcenter.com/cearticle?an=00006247-201510000-00007&Journal_ID=54013&Issue_ID=3208683).
- Redacted. (2018). Facts and Figures. Retrieved from

<https://annualreports.vcu.edu/vcuhealth/facts.html>.

Redacted. (2019). Mission and Vision. Retrieved from <https://www.vcuhealth.org/our-story/who-we-are/mission-and-vision>.

Saddawi-Konekfa, D., Schumacher, D. J., Baker, K. H., Charmin, J. E., & Gollwitzer, P. M.

(2016). Changing Physician Behavior with Implementation Intentions: Closing the Gap Between Intention and Actions. *Academic Medicine: Journal of the Association of American Medical Colleges*, 91(9), 1211-1216. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/27008360>.

Schuller, K. A., Lin, S., Gamm, L., & Edwardson, N. (2015). Discharge Phone Calls: A Technique to Improve Patient Care During the transition From Hospital to Home. *The Journal for Healthcare Quality*, 37(3) 163-172. Retrieved from [https://journals.lww.com/jhqonline/Fulltext/2015/05000/Discharge\\_Phone\\_Calls\\_\\_\\_A\\_Technique\\_to\\_Improve.2.aspx](https://journals.lww.com/jhqonline/Fulltext/2015/05000/Discharge_Phone_Calls___A_Technique_to_Improve.2.aspx).

Sorita, A., Robelia, P. M., Kattel, S. B., McCoy, C. P., Keller, A. S., Almars, J., ... Kashiwagi, D. T. (2017). The Ideal Hospital Discharge Summary: A Survey of U.S. Physicians. *Journal of Patient Satisfaction*, 00(00). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/28885382>.

Stossel, L. M., Segar, N., Gliatto, P., Fallar, R., & Karani, R. (2012). Readability of Patient Education Materials Available at the Point of Care. *Journal of General Internal Medicine*, 27(9), 1165-1170. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3514986/>.

Terry, J., & Cutter, J. (2013). Does Education Improve Mental Health Practitioners' Confidence

- in Meeting the Physical Health Needs of Mental Health Service Users? A Mixed Methods Pilot Study. *Issues in Mental Health Nursing*, 34(4), 249-255. Retrieved from <https://www.tandfonline.com/doi/abs/10.3109/01612840.2012.740768>.
- The Joint Commission. (2007). "What Did the Doctor Say?:" Improving Health Literacy to Protect Patient Safety. Retrieved from [https://www.jointcommission.org/assets/1/18/improving\\_health\\_literacy.pdf](https://www.jointcommission.org/assets/1/18/improving_health_literacy.pdf).
- Titler, M. G. (2008). Patient Safety and Quality: An Evidence-Based Handbook for Nurses. *Agency for Healthcare Research and Quality*. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK2659/>.
- Unaka, N. I., Statile, A., Haney, J., Beck, A. F., Brady, P. W., Jerardi, K. E. (2017). Assessment of readability, understandability, and completeness of pediatric hospital medicine discharge instructions. *Journal of Hospital Medicine*, 12(2), 98-101. Retrieved from <https://europepmc.org/abstract/med/28182805>.
- U.S. Department of Health and Human Services & Centers for Disease Prevention and Control. (2009). Simply Put, a guide for creating easy-to-understand materials. Retrieved from [https://www.cdc.gov/healthliteracy/pdf/simple\\_put.pdf](https://www.cdc.gov/healthliteracy/pdf/simple_put.pdf).
- U.S. Department of Health and Human Services. (2000). Healthy People 2010: understanding and improving health, 2nd ed. Washington, DC: U.S. Government Printing Office; November 2000.
- Waugaman, W. R., & Lohrer, D. J. (2000). From nurse to nurse anesthetist: The influence of age and gender on professional socialization and career commitment of advanced practice nurses. *Journal of Professional Nursing*, 16(1), 47-56. Retrieved from <https://www.sciencedirect.com/science/article/abs/pii/S8755722300800113>



- Waniga, H. M., Gerke, T., Shoemaker, A., Bourgoine, D., & Eamranond, P. (2016). The Impact of Revised Discharge Instructions on Patient Satisfaction. *Journal of Patient Experience*, 3(3), 64-68. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5513645/>.
- Young, L. J., & Newell, K. J. (2008). Can a Clinical Continuing Education Course Change Behavior in Dental Hygiene Practice? *Journal of Dental Hygiene*, 82(4). Retrieved from <http://jdh.adha.org/content/jdentyg/82/4/33.full.pdf>.